

IMMEDIATE DENTURE



CONTENTS:

DEFINITION TYPES OF IMMEDIATE DENTURE **INDICATION** CONTRAINDICATION DIAGNOSIS TREATMENT PLANNING **ADVANTAGES** DISADVANTAGES **CLINICAL PROCEDURE**

IMMEDIATE DENTURE

An immediate denture is "a complete denture or removable partial denture fabricated for placement immediately after the removal of natural teeth".

- Immediate dentures may be either single immediate dentures or upper and lower immediate dentures in the same patient
- Preferably one denture
- An immediate denture can also be an overdenture

INDICATIONS FOR IMMEDIATE DENTURE

- Esthetics.
- Periodontally weak tooth.





CONTRAINDICATION OF IMMEDIATE DENTURE

- Patients who are in poor general health or who are poor surgical risks.
- Patients who are identified as uncooperative






TYPES OF IMMEDIATE DENTURE:

- 1.Conventional or classic
- 2.Interim or transitional

CONVENTIONAL IMMEDIATE DENTURE

- Immediate denturemodified as permanent denture
- Two surgical appointments
- Extraction of posterior except I premolar
- 6- 8 weeks allow ridges to heal followed by anterior removal
- Immediate denture insertion



INTERIM IMMEDIATE DENTURE

Used when anterior and posterior teeth remain and placement of the immediate denture after the extraction.



DIAGNOSIS, TREATMENT PLANNING

1.ORAL EXAMINATION:

- Periodontal probings.
- Full charting of the teeth.
- Frenum.
- Tori reduction if necessary.
- Evaluation and palpation of potential denture supporting tissues and PPS.











2.ORAL PROPYLAXIS:

- Scaling –remove the calculus
- Reduce postoperative oedema and infection.




3.TOOTH MODIFICATION:

- Caries removal and restoration.
- Periodontal diseases elimination.
- Elimination of occlusal discripancies.
- occlusal modification.



4.DIAGNOSTIC CAST MOUNTING:

- Evaluation of tooth position.
- Jaw relationships.
- Occlusal plane discripancies.
- Analyse the undercut




5.RADIOGRAPHIC EXAMINATION:

- Extent of bone loss.
- Root pieces.
- Impacted tooth.
- Foreign bodies.
- Exostosis.
- Cyst, tumors etc....

CLINICAL PROCEDURES:

CONVENTIONAL TYPE


































INTERIM TYPE











Before

After

ADVANTAGES OF IMMEDIATE DENTURE

- Patient perspection
- No time patient without denture
- Less apprehension
- To continue social and business activities
- Uninterupted digestive function
- Get adjusted to changes more quickly
- Appearance
- Speech and chewing habits

ADVANTAGES OF IMMEDIATE DENTURE

- Maintenance the Circumoral support
- ID acts as a bandage over the surgical site, controls haemorrhage, prevents contamination, protects wound.
- Less postoperative pain
- Reduce residual ridge resorption.
- Ridge preservation
- size, shape, shade ,inclination of natural teeth can be duplicated





DISADVANTAGES OF IMMEDIATE DENTURE

- Increased treatment complexity
- Try- in is not possible
- Increased treatment cost.
- Increased patient visits.
- Fit of denture

PATIENT INSTRUCTION

- Advised for post-extraction precautions and discharged.
- Warm salt water rinses.
- Avoid mouthwashes containing alcohol
- Not to remove the denture for 24 hrs.
- After 24hrs patient was recalled prosthesis was removed, checked & cleaned.
- Patient recalled after 1 week for review.

